

WebM&M

Morbidity and Mortality Rounds on the Web

Spotlight

Delayed Diagnosis and Treatment of Systemic Lupus Erythematosus with a Psychiatric Presentation



Agency for Healthcare Research and Quality
Advancing Excellence in Health Care



Source and Credits

- This presentation is based on the March 2024 AHRQ WebM&M Spotlight Case
 - See the full article at <https://psnet.ahrq.gov/webmm>
 - CME credit is available
 - Editor's Note: this case was adapted from Capozzola DD, Terrence J. *Failure to Diagnose and Treat Lupus May Support Liability for Psychiatrist*. Healthcare Risk Management (Relias Media). Published August 1, 2022.
- Commentary by: James A. Bourgeois, OD, MD and Glen Xiong, MD
- AHRQ WebM&M Editors in Chief: Patrick Romano, MD, MPH and Deb Bakerjian, PhD, APRN, RN
 - Spotlight Editors: Patrick Romano, MD, MPH and Noelle Boctor, MD
 - Managing Editor: Meghan Weyrich, MPH

Objectives

At the conclusion of this educational activity, participants should be able to:

- Describe the clinical manifestations of a primarily psychiatric presentation of systemic lupus erythematosus (SLE).
- Identify elements of the patient history that raise the probability of SLE in atypical psychotic disorder.
- Describe atypical features of primary psychotic illness that raise suspicion of SLE.
- Discuss the need for clear communication about clinical data between medical specialists.
- Appreciate that implicit bias (e.g., society stigma) about psychiatric illness may impact the degree of suspicion for psychiatric presentation of systemic illness.

DELAYED DIAGNOSIS AND TREATMENT OF SYSTEMIC LUPUS ERYTHEMATOSUS WITH A PSYCHIATRIC PRESENTATION

A case highlighting the clinical manifestations of a primarily psychiatric presentation of systemic lupus erythematosus (SLE), the importance of family history when evaluating patients with psychotic presentations, and the need for clear communication between medical specialists to ensure safe, high-quality care

Case Details (1)

- An 18-year-old woman with no significant past medical history was admitted to a community hospital from a behavioral health center for evaluation and treatment of acute psychosis with paranoid delusions.
- The nurse practitioner managing the patient was uncertain as to whether the psychosis was due to a “primary” psychiatric condition (e.g., schizophreniform disorder, schizophrenia) or a systemic medical condition, so two consultations were obtained.
- On hospital day 5, a psychiatrist evaluated the patient and recommended initiation of an antipsychotic medication.

Case Details (2)

- On hospital day 7, the nurse practitioner learned from the patient's father that there was a family history of systemic lupus erythematosus (SLE) and suggested that the patient be evaluated for lupus.
- Laboratory tests on hospital day 9 indicated mild pancytopenia (e.g., platelet count 120,000/microliter), an elevated antinuclear antibody (ANA), and abnormally elevated anti-double-stranded DNA (deoxyribonucleic acid) (anti-ds DNA).
- Urinalysis, physical examination, and electrocardiogram were normal.
- The patient was discharged on hospital 10, but none of the treating physicians reviewed the test results until 2-3 days after discharge.
- The patient was never referred for further evaluation of these findings.

Case Details (3)

- The patient's delusions and other psychotic symptoms abated over the following months, but she was admitted to a second hospital for exacerbation of psychotic symptoms six months after her first hospitalization.
- The medical team at this hospital quickly discovered the abnormal test results from six months earlier, documented the diagnosis of SLE, and initiated appropriate corticosteroid treatment.
- The patient ultimately died of multiple organ failure.

DELAYED DIAGNOSIS AND TREATMENT OF SYSTEMIC LUPUS ERYTHEMATOSUS WITH A PSYCHIATRIC PRESENTATION

THE COMMENTARY

By James A. Bourgeois, OD, MD and Glen Xiong, MD

BACKGROUND

Background (1)

- This unfortunate case likely represented a presentation of psychotic illness as *forme fruste* or prodrome of systemic lupus erythematosus (SLE).
 - Psychiatric prodromal states, wherein an initial “psychiatric” presentation heralds an *ultimate* course of relapsing, multisystem involvement, represent an accepted construct for multiple sclerosis and an emerging construct for SLE, based on recent research.^{1,2,3}
 - This prodromal construct remains somewhat controversial, especially when the prodromal state primarily affects the central nervous system (CNS) and the full syndrome (as in SLE) more dramatically affects other organ systems.
 - In SLE, the “primarily rheumatologic” full syndrome is not always clearly connected to a past psychiatric prodrome, and the psychiatric symptoms that patients experience when their multisystem illness is diagnosed can be attributed to a maladaptive emotional reaction to the diagnosis, rather than to CNS manifestations of SLE itself.

Background (2)

- This is essentially a “lumper/splitter” argument.
 - If one accepts that SLE is a centrally mediated, multisystem illness, it is reasonable to understand a psychiatric prodrome as heralding the initial onset of SLE with later, concurrent multisystem and CNS involvement attributable to the same underlying pathophysiology.
 - The psychiatric prodromal concept also may lack predictive value, in that many patients (eventually) diagnosed with SLE or multiple sclerosis *may not*, in retrospect, have had a clear psychiatric prodrome.
- This case exemplifies the classic “zebra” in clinical diagnosis.
 - Given the high prevalence of primary psychiatric conditions in young adults, it is difficult to estimate the posterior probability or likelihood of rare conditions that present with initial psychiatric symptoms in this population.
 - Therefore, diagnostic and treatment approaches generally rely on expert clinical experience attuned to subtle atypical features rather than diagnostic protocols based on randomized controlled trials.

DIAGNOSING PSYCHOTIC ILLNESS

Diagnosing Psychotic Illness (1)

- Psychotic illness that is *ultimately* attributed to schizophrenia, other psychotic disorders, and other “primary” psychiatric illnesses with psychotic features is far more common in the young adult population than a psychotic presentation of systemic illness.
 - The differential diagnosis for psychiatric illnesses presenting as acute psychosis in young adult patients includes:
 - Schizophreniform disorder
 - Schizophrenia
 - Brief psychotic disorder
 - Major depressive disorder with psychotic features
 - Bipolar disorder (major depressive or manic episode)
 - Substance-induced psychotic disorder
 - Autistic disorder
 - Intellectual disability
 - Posttraumatic stress disorder
 - Borderline personality disorder

However, these diagnoses are established only after health professionals use the medical history, physical examination, laboratory testing, and imaging studies (if appropriate) to effectively rule out other causes of psychotic illness.

Diagnosing Psychotic Illness (2)

An important concept in the diagnosis of psychotic illness is “atypical features”, referring to psychotic symptoms that are not typical for schizophrenia or other “primary psychotic” illnesses.

Psychotic presentations with *atypical features* lead to greater suspicion of a systemic, rather than “purely psychiatric,” illness, but this combination of findings also lacks specificity in many cases. Presentations with atypical features should lead to a more thorough diagnostic work-up, potentially including neuroimaging, electroencephalography [EEG], robust laboratory assessment, and (only in selected cases) cerebrospinal fluid [CSF] studies.⁴

Psychotic Presentations with Atypical Features:

- Visual, olfactory, and/or tactile rather than auditory hallucinations
- Prominent cognitive impairment
- Concurrent motor signs
- Catatonia
- Atypical age of onset
- Confusion
- Disorientation
- Language disturbance
- Misidentification delusions (Capgras syndrome)
- Constitutional symptoms (e.g., significant weight loss)
- Lack of family history of psychotic disorders
- Lack of personal history of schizoid personality disorder

Diagnosing Psychotic Illness (3)

- Neuropsychiatric SLE (NPSLE) is reported in 25-75% of SLE patients and may include cognitive impairment, headache, anxiety, depressed mood, psychosis, seizures, sleep disturbances, and abnormal movements.⁴⁻⁷
- This constellation of *multiple* CNS symptoms comorbid with psychosis, especially in a young patient with new-onset psychotic illness who may have a family history of SLE or other autoimmune disease, should increase the suspicion of NPSLE as the cause of first-episode psychosis.⁴⁻⁷
- Mechanistically, autoimmunity may be associated with antibodies to synaptic and neuronal cell membrane proteins (e.g., anti-N-methyl-d-aspartate receptor antibodies).⁴

Diagnosing Psychotic Illness (4)

- This complex presentation of multiple CNS findings in addition to psychosis would not be typical of primary psychotic illness (e.g., schizophrenia, schizophreniform disorder).
- NPSLE can also be classified into diffuse syndromes associated with the CNS (as above) and focal syndromes associated both with the CNS (e.g., cerebrovascular disease) and the peripheral nervous system (e.g., Guillain-Barre syndrome, myasthenia gravis).⁶
- NPSLE risk appears to be higher in early-onset SLE than late-onset SLE.^{5,8}
- NPSLE symptoms may evolve over time, with systemic findings such as inflammatory skin lesions and hematologic abnormalities appearing later.³

Diagnosing Psychotic Illness (5)

- The diagnostic challenge remains regarding how to work up first presentation of psychotic illness to assure that systemic illnesses presenting as a psychotic episode are fully considered, as clinical intervention is different for psychotic illnesses attributable to systemic illness.
- A *brief* list of other illnesses that can present as isolated psychosis include: ⁹⁻¹⁴
 - Substance-related conditions;
 - Syndrome of inappropriate secretion of antidiuretic hormone (SIADH) with hyponatremia;
 - Liver disease (such as hepatitis B or C with cirrhosis and hepatic encephalopathy);
 - Hyper- or hypothyroidism;
 - Folate, vitamin B12, or vitamin D deficiency;
 - Rheumatologic diseases;
 - Wilson's disease; human immunodeficiency virus (HIV) disease;
 - Neurosyphilis.

Diagnosing Psychotic Illness (6)

- The physical examination in this setting should include a full neurological exam, assessment of height and weight, and vital signs.⁹
- It is common and reasonable, although relatively low yield, to obtain unenhanced computed tomography (CT) of the head, but not contrast-enhanced CT or magnetic resonance imaging (MRI) for a first presentation of psychotic illness.
 - In *established* psychotic illness, neuroimaging is not repeated at each exacerbation, particularly if the patient experiences stereotypical recurrences.

Diagnosing Psychotic Illness (7)

Several laboratory studies are routinely performed in this setting, as vitamin deficiencies and other metabolic and endocrine abnormalities can present with psychotic symptoms.

Laboratory Studies for First Episode Psychotic Illness⁹⁻¹⁴

- Urine drug screen
- Basic metabolic panel
- Liver associated enzymes
- Thyroid stimulating hormone
- Vitamin B12
- Vitamin D
- Folate
- Antinuclear antibody (ANA) screen
- Ceruloplasmin
- Erythrocyte sedimentation rate (ESR)
- Human immunodeficiency virus
- Antitreponemal IgG
- Cerebrospinal Fluid analysis (ESR, cell count, and, when indicated, antineoplastic antibodies, limbic encephalitis antibody panels)

Other laboratory studies can be ordered, depending on the clinical context of presentation.^{9,10} More invasive laboratory studies, such as lumbar puncture for cerebrospinal fluid (CSF) studies, is not routinely accomplished initially. While SLE with a psychotic prodrome is relatively well-known but uncommon, routine testing for SLE-specific rheumatologic markers is not routine, nor is assessment for anti-NMDA-receptor encephalitis (ANMDARE) on initial evaluation. As SLE is highly heritable, once the family history of SLE was ascertained, the nurse practitioner appropriately obtained SLE-specific laboratory studies, with positive results.^{7,8}

Diagnosing Psychotic Illness (8)

- The combination of a psychotic presentation with positive SLE-specific laboratory results is enough to conclude that this patient probably had SLE-related psychotic disorder, but the patient should then have promptly had CSF studies, MRI, and an urgent rheumatology consult for a trial of immunomodulation.¹
 - In such a case, immunomodulation (often with corticosteroids) may improve psychotic symptoms. Ironically, corticosteroids *themselves* (especially high dose prednisone) may improve the systemic manifestations but worsen the CNS disease, rendering the patient “systemically improved but psychiatrically worse”, at least temporarily.¹⁵
 - Other immunomodulators may also be used in NPSLE just as in systemic SLE. High quality evidence for the optimal treatment approach is lacking.¹⁶
 - Smoking may be associated with increased SLE disease activity, so smoking cessation is strongly encouraged.^{3,7}

Diagnosing Psychotic Illness (9)

- An empiric trial of antipsychotic pharmacotherapy would be typical in a case of this type, even with concurrent immunomodulation, though there would be uncertainty about whether any subsequent clinical improvement was attributable to the antipsychotic, immunomodulator, or both.
- If a patient with SLE-related psychotic illness has a robust treatment response, they may not need the several months of antipsychotic prophylaxis that would be routine in purely psychotic illness (e.g., schizophreniform disorder, schizophrenia).
- Antipsychotic treatment resistance is a common feature and should lead clinicians to look for additional CNS or systemic causes of new-onset psychosis.

Diagnosing Psychotic Illness (10)

- This patient did not demonstrate antipsychotic treatment resistance, providing another missed opportunity for timely diagnosis.
 - More specifically, the CNS SLE abated with antipsychotic treatment, even without immunomodulation.
 - There was apparently no other manifestation of CNS SLE such as CNS inflammation or systemic symptoms such as headache, cognitive impairment, seizures, and stroke.
- In the second clinical encounter, it appeared that a psychotic exacerbation occurred, but again systemic symptoms were not readily apparent. Despite the possibility of targeted treatment, the patient died suddenly of organ failure.

APPROACHES TO IMPROVING PATIENT SAFETY

Approaches to Improving Patient Safety (1)

- The tragedy in this case is that there was clinical suspicion of SLE based on the family history, and at least initially appropriate screening laboratory studies were done and reported as positive.
- This profile would have obligated an urgent rheumatology referral and inpatient admission for CSF studies, MRI, and trial of immunomodulation.
- Close collaboration with consultation-liaison psychiatry would have been needed to manage antipsychotic medication, monitor for and manage delirium, and adjust psychotropic medications depending on the patient's response.
- This patient could have had a favorable result with this management plan, but we will never know with certainty. It is unclear why obviously abnormal laboratory studies were not acted upon to effectuate the needed rheumatology consult and urgent SLE work up.

Approaches to Improving Patient Safety (2)

- Failure to follow-up on abnormal laboratory results is not an uncommon system or process error and can lead to dire consequences for patients.
- Automatic checks may be implemented in electronic health record (EHR) systems to ensure that laboratory results are reviewed (and marked as reviewed).
- If a critical laboratory result has not been reviewed within a certain time period, automatic prompts may be sent to the ordering physician as well as the primary physician of record.

Approaches to Improving Patient Safety (3)

- In this case, the lack of apparent systemic symptoms and predominant presentation of psychosis reduced the likelihood of the patient seeking primary medical care.
- Finally, psychotic and cognitive symptoms may have also impaired the patient's ability to appropriately perceive subtle systemic symptoms.
- In this context, the lack of a primary care physician to coordinate post-discharge care may have contributed to the failure to follow up on abnormal laboratory tests.

TAKE HOME POINTS

Take-Home Points

- Systemic lupus erythematosus (SLE) involving the central nervous system can have a primarily psychotic presentation at illness onset.
- Psychotic presentation may be a prodrome for later presentation of full spectrum SLE.
- Psychotic presentations in patients with strong family history or personal history of SLE mandate appropriate laboratory screening and other interventions to rule in or rule out the diagnosis of SLE.
- Psychotic symptoms, even if attributable to SLE, can still respond to antipsychotic treatment but such patients still need full systemic disease workup and monitoring.
- Health systems should develop processes to facilitate timely follow-up on ordered tests with pending results, even after discharge.

REFERENCES

References

1. Govoni M, Bortoluzzi A, Padovan M, et al. The diagnosis and clinical management of the neuropsychiatric manifestations of lupus. *J Autoimmun.* 2016;74:41-72. [[Free full text](#)]
2. Wijnands JM, Zhu F, Kingwell E, et al. Five years before multiple sclerosis onset: Phenotyping the prodrome. *Mult Scler.* 2019;25(8):1092-1101. [[Available at](#)]
3. Frazzei G, van Vollenhoven RF, de Jong BA, Siegelaar SE, van Schaardenburg D. Preclinical Autoimmune Disease: a Comparison of Rheumatoid Arthritis, Systemic Lupus Erythematosus, Multiple Sclerosis and Type 1 Diabetes. *Front Immunol.* 2022;13:899372. [[Free full text](#)]
4. Najjar S, Steiner J, Najjar A, Bechter K. A clinical approach to new-onset psychosis associated with immune dysregulation: the concept of autoimmune psychosis. *J Neuroinflammation.* 2018;15(1):40. [[Free full text](#)]
5. van der Heijden H, Rameh V, Golden E, et al. Implications of Inflammatory Processes on a Developing Central Nervous System in Childhood-Onset Systemic Lupus Erythematosus. *Arthritis Rheumatol.* Published online October 30, 2023. [[Available at](#)]
6. Emerson JS, Gruenewald SM, Gomes L, Lin MW, Swaminathan S. The conundrum of neuropsychiatric systemic lupus erythematosus: Current and novel approaches to diagnosis. *Front Neurol.* 2023;14:1111769. [[Free full text](#)]
7. Mizrahi M, Anderson E, Carroll KR, Tehrani N, Volpe BT, Diamond B. Cognitive dysfunction in SLE: An understudied clinical manifestation. *J Autoimmun.* 2022;132:102911. [[Available at](#)]
8. Pamuk ON, Raza AA, Hasni S. Neuropsychiatric lupus in late- and early-onset systemic lupus erythematosus patients: a systematic review and meta-analysis. *Rheumatology (Oxford).* 2024;63(1):8-15. [[Available at](#)]
9. Dorney K, Murphy M. Recommendations for the medical work-up of first episode psychosis, including specific relevance to Indigenous Australians: A narrative review. *Early Interv Psychiatry.* 2021;15(3):423-438. [[Available at](#)]
10. Skikic M, Arriola JA. First Episode Psychosis Medical Workup: Evidence-Informed Recommendations and Introduction to a Clinically Guided Approach. *Child Adolesc Psychiatr Clin N Am.* 2020;29(1):15-28. [[Available at](#)]
11. Esque J, Rasmussen A, Spada M, Gopalan P, Sarpal D. First-Episode Psychosis and the Role of the Psychiatric Consultant. *J Acad Consult Liaison Psychiatry.* 2022;63(1):32-35. [[Available at](#)]
12. Pollak TA, Rogers JP, Nagele RG, et al. Antibodies in the Diagnosis, Prognosis, and Prediction of Psychotic Disorders. *Schizophr Bull.* 2019;45(1):233-246. [[Free full text](#)]
13. Chapman MR, Vause HE. Anti-NMDA receptor encephalitis: diagnosis, psychiatric presentation, and treatment. *Am J Psychiatry.* 2011;168(3):245-251. [[Available at](#)]
14. Kayser MS, Kohler CG, Dalmau J. Psychiatric manifestations of paraneoplastic disorders. *Am J Psychiatry.* 2010;167(9):1039-1050. [[Available at](#)]
15. Dubovsky AN, Arvikar S, Stern TA, Axelrod L. The neuropsychiatric complications of glucocorticoid use: steroid psychosis revisited. *Psychosomatics.* 2012;53(2):103-115. [[Available at](#)]
16. Schwartz N, Stock AD, Putterman C. Neuropsychiatric lupus: new mechanistic insights and future treatment directions. *Nat Rev Rheumatol.* 2019;15(3):137-152. [[Free full text](#)]